

Premier Health Plan

POLICY AND PROCEDURE MANUAL

Policy Number: MP.114.PH
Last Review Date: 02/08/2018
Effective Date: 04/01/2018

MP.114.PH – High-Resolution Anoscopy

This policy applies to the following lines of business:

- ✓ Premier Employee

Premier Health Plan considers **High-Resolution Anoscopy (HRA)** medically necessary for members Referred for HRA that have both of the following criteria:

1. Part of patient population at increased risk for anal cancer including any of the following:
 - a. Men having sex with men (MSM); or
 - b. Men and women with HIV disease; or
 - c. Women with a history of high-grade genital dysplasia (cervical, vaginal and vulvar); or
 - d. HPV patients especially those with a history of genital warts, either internal or external; or
 - e. Solid organ transplant recipients who are immunosuppressed; or
 - f. Long-term corticosteroid users; or
 - g. Smokers
2. Anal cytology findings of any of the following:
 - a. ASC-US (atypical squamous cells of undetermined significance) = AIN I; or
 - or
 - b. LSIL (low-grade squamous intraepithelial lesion) = AIN I; or
 - c. ASC-H (atypical squamous cell, cannot rule out a high grade lesion) = AIN II; or
 - d. HSIL (high-grade squamous intraepithelial lesion) = AIN II or III

HRA referrals for anal symptoms suspicious of dysplastic progression in which anal cytopathology is not available will be reviewed on a case-by-case basis.

These include members with either of the following conditions:

1. Solid organ transplant candidates who are immunosuppressed
2. Women with high grade genital dysplasias or history of vulvar and cervical cancer

Frequency of follow-up with HRA generally includes the following:

1. Normal findings – repeat cytology in 1 year
2. ASC-US, LSIL, ASC-H, or HSIL
 - Patients with AIN I can be followed up every 6 -12 months
 - Patients with AIN II or III - therapy is recommended with follow-up in 6 months post therapy

MP.114.PH – High-Resolution Anoscopy

Policy Number: MP.114.PH
Last Review Date: 02/08/2018
Effective Date: 04/01/2018

Limitations

- HRA is **not covered** for routine screening (only indicated for diagnostic use – after abnormality detected during screening)
- Coverage of this procedure is limited to physicians or advanced practice clinicians who have completed comprehensive training in HRA such as provided through the ASCCP/AMC (AIDS Malignancy Consortium)/ACTG (Adult AIDS Clinical Trials Group) High Resolution Anoscopy (HRA) certification process.
- Coverage of HRA, when performed in conjunction with treatment/destruction of the anal dysplastic lesions, will be considered global to the primary procedure.

Background

Anal dysplasia, caused by the human papillomavirus (HPV), is defined as abnormal cells or lesions in the lining of the anal canal. Although the incidence of anal cancer is low in the United States, if it is detected early it can be treated successfully. The incidence varies depending on the presence of risk factors such as multiple sex partners, HPV and/or HIV infection, receptive anal intercourse, history of anal warts, STIs and/or fissures, being over 50 years old, women with a history of cervical cancer, and smoking cigarettes.

High-resolution anoscopy (HRA) is a minimally invasive procedure for more detailed identification, management and treatment of anal dysplasia in high-risk populations. During the HRA procedure, a lubricated anoscope is inserted into the anal canal. A cotton swab wrapped in gauze and soaked in 3-percent acetic acid is then inserted through the anoscope, and the anoscope is removed, leaving the gauze in place. The acetic acid gives dysplastic epithelium a white appearance. After two minutes, the gauze is removed and the anoscope reinserted. A high-resolution colposcope (magnification of 10x to 40x) is used to view the walls of the anus. A biopsy of suspicious tissue can be taken. The procedure is generally performed in an office setting in either a bent over or lying position and usually takes approximately 15 minutes.

Codes:

| CPT Codes / HCPCS Codes / ICD-10 Codes | |
|--|---|
| Code | Description |
| CPT Codes | |
| 46601 | Anoscopy; diagnostic, with high-resolution magnification (HRA) (eg, colposcope, operating microscope) and chemical agent enhancement, |

MP.114.PH – High-Resolution Anoscopy

Policy Number: MP.114.PH
 Last Review Date: 02/08/2018
 Effective Date: 04/01/2018

| | |
|--|---|
| | including collection of specimen(s) by brushing or washing, when performed |
| 44607 | Anoscopy; with high-resolution magnification (HRA) (eg, colposcope, operating microscope) and chemical agent enhancement, with biopsy, single or multiple |
| ICD-10 codes covered if selection criteria are met: | |
| A63.0 | Anogenital (venereal) warts |
| B20 | Human immunodeficiency virus (HIV) disease |
| B97.35 | Human immunodeficiency virus, type 2 (HIV 2) |
| B97.7 | Papillomavirus as the cause of diseases classified elsewhere |
| C20-C21.8 | Malignant neoplasm of rectum, anal canal, and anus |
| D12.8-D12.9 | Benign neoplasm of rectum, anus, and anal canal |
| D01.3 | Carcinoma in situ of anus and anal canal |
| K62.0-K62.1 | Anal and rectal polyp |
| K62.5 | Hemorrhage of anus and rectum |
| K62.6 | Ulcer of anus and rectum |
| K62.81 | Anal Sphincter tear (healed) (nontraumatic) (old) |
| K62.82 | Dysplasia of anus |
| K62.89 | Other specified diseases of anus and rectum |
| N87.0-N87.9 | Dysplasia of cervix |
| N89.0-N89.3 | Dysplasia of vagina |
| R85.6-R85.619 | Abnormal cytologic smear of anus |
| R85.81-R85.82 | Anal high-low risk human papillomavirus (HPV) DNA test positive |
| Z21 | Asymptomatic human immunodeficiency virus (HIV) infection status |
| Z72.52 | High risk homosexual behavior |
| Z72.53 | High risk bisexual behavior |
| Z79.51-Z79.52 | Long-term use (current) of steroids |
| Z87.410 | Personal history of cervical dysplasia |
| Z87.411 | Personal history of vaginal dysplasia |
| Z87.412 | Personal history of vulvar dysplasia |
| Z94.0-Z94.9 | Organ or tissue replaced by transplant |

MP.114.PH – High-Resolution Anoscopy

Policy Number: MP.114.PH
Last Review Date: 02/08/2018
Effective Date: 04/01/2018

| | |
|-------|-----------------------------------|
| Z95.3 | Presence of xenogenic heart valve |
|-------|-----------------------------------|

References

1. Boston Medical Center Obstetrics & Gynecology. Anal Dysplasia and Anal Cancer. <https://www.bmc.org/obgyn/services/specialprograms-anal-cancer-dysplasia.htm>
2. Centers of Medicare and Medicaid Services (CMS). Local coverage article: Coverage for High Resolution Anoscopy A53408. (Palmetto GBA): Revision Effective Date March 16, 2017. <https://www.cms.gov/medicare-coverage-database/details/article-details.aspx?articleId=53408&ver=10&Date=&DocID=A53408&bc=hAAAABAAA&AAA&AAA&>
3. Department of Health & Human Services (HHS). Agency for Healthcare Research and Quality (AHRQ). National Guideline Clearinghouse (NGC). Screening for Cervical Cancer NGC-9372. Last Updated: December 14, 2012. <http://www.guideline.gov/content.aspx?id=38572>
4. Gimenez F., Costa-e-Silva,IT, Daumas A, et al. The value of high-resolutions anoscopy in the diagnosis of anal cancer precursor lesions in HIV-positive patients. Arq Gastroenterol. 2011 Jan; 48(2):136-145. <http://www.scielo.br/pdf/ag/v48n2/a10v48n2.pdf>
5. Hayes Medical Technology Directory. High-Resolution Anoscopy for the Evaluation of Anal Lesions. Annual Review October 22, 2015.
6. Steele SR, Varma MG, Melton GB, et al. Practice parameters for anal squamous neoplasms (Standards Practice Task Force of the American Society of Colon and Rectal Surgeons). Dis Colon Rectum. 2012 Jul; 55(7): 735-749. https://www.fascrs.org/sites/default/files/downloads/publication/practice_parameters_for_anal_squamous_neoplasms.21.pdf
7. U.S. Department of Health & Human Services (HHS). Agency for Healthcare Research and Quality (AHRQ). National Guideline Clearinghouse (NGC). Neoplastic complications of HIV infection. NGC- 5840. Issued: September 20, 2007. <http://f.i-md.com/medinfo/material/d4a/4eb2251e44ae9d3f78b07d4a/4eb2253844ae9d3f78b07d4d.pdf>
8. U.S. Department of Health & Human Services (HHS): Health Resources and Services Administration. HIV/AIDS Bureau: Guide for HIV/AIDS Clinical Care - Section 6: Comorbidities and Complications- Anal Dysplasia. April 2014. <http://hab.hrsa.gov/deliverhivaidscares/2014guide.pdf>
9. U.S. Department of Health & Human Services (HHS). Agency for Health Research and Quality (AHRQ): National Guideline Clearinghouse (NGC):

MP.114.PH – High-Resolution Anoscopy

Policy Number: MP.114.PH
Last Review Date: 02/08/2018
Effective Date: 04/01/2018

- Primary care approach to the HIV-infected patient. NGC-8648- Last updated: October 27, 2011. <http://www.guideline.gov/content.aspx?id=34268>
10. United States Department of Veterans Affairs: VA National HIV/AIDS Website. Anal dysplasia. Primary care of veterans with HIV. Updated October 28, 2011. <http://www.hiv.va.gov/provider/manual-primary-care/anal-dysplasia.asp>

Disclaimer:

Premier Health Plan medical payment and prior authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. The policies constitute only the reimbursement and coverage guidelines of Premier Health Plan and its affiliated managed care entities. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies.

Premier Health Plan reserves the right to review and update the medical payment and prior authorization guidelines in its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

These policies are the proprietary information of Evolent Health. Any sale, copying, or dissemination of said policies is prohibited.